

LEBANON Scape®

STARTER FERTILIZER
WITH 0.08% MESO
PREEMERGENT WEED CONTROL

21-22-4

GUARANTEED ANALYSIS

Total Nitrogen (N)	21%
4.5% Ammoniacal Nitrogen	
9.1% Urea Nitrogen	
6.8% Other Water Soluble Nitrogen**	
0.6% Other Water Insoluble Nitrogen**	
Available Phosphate (P₂O₅)	22%
Soluble Potash (K₂O)	4%
Sulfur (S)	1%
1.0% Combined Sulfur (S)	
Iron (Fe).....	0.5%
0.5% Water Soluble Iron (Fe)	
Derived From: Methyleneurea, Urea, Potassium Sulfate, Ammonium Phosphate and Ferrrous Sulfate.	
** Contains 5.6% slowly available nitrogen from methyleneurea.	

F689

ACTIVE INGREDIENTS:	EPA Reg. No. 538-377-981
Mesotrifluron	0.08%
Other Inert Ingredients	99.92%
TOTAL	100.00%
TCAS #104208-502-8	

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS AND DOMESTIC ANIMALS
 CAUTION: Causes moderate eye irritation. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum, using tobacco or using the toilet.

FIRST AID

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor immediately for treatment advice. Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. Applicator and other handlers must wear: Long-sleeved shirt, long pants, socks and shoes, and chemical resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride.

User Safety Requirements:

Follow manufacturer's instructions for cleaning/maintaining Personal Protective Equipment. If there are no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:
 • Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
 • Remove clothing/PPE immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
 • Remove PPE immediately after handling this product. Wash outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

ENVIRONMENTAL HAZARDS

Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not contaminate water when disposing of equipment wash waters or rinsate.

Surface Water Advisory

This product may contaminate water through drift of spray in wind. This product has a high potential for runoff for several weeks after application. Poorly draining soils and soils with shallow water tables are more prone to produce runoff that contains this product. A leach well maintained vegetative buffer strip between areas to which this product is applied and surface water features such as ponds, streams, and springs will reduce the potential for contamination of water from runoff. Runoff of this product will be reduced by avoiding applications when rainfall is forecasted to occur within 48 hours. Sound erosion control practices will reduce this product's contribution to surface water contamination.

DIRECTIONS FOR USE

It is a violation to use this product in manner inconsistent with its labeling. Do not apply this product in a way that the contents or other parts thereof, either directly or through drift, only protected handlers may be in the area during application; for any requirements specific to your State or Tribe, contact the agency responsible for pesticide regulation.

Applicational Use Requirements

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR Part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on the label about Personal Protective Equipment (PPE) and restricted entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

DO NOT ENTER OR ALLOW WORKERS INTO TREATED AREAS DURING THE RESTRICTED ENTRY INTERVAL (REI) OF 12 HOURS. Except for the REI, this product is not a restricted entry pesticide as defined by the Worker Protection Standard, under certain circumstances, allow workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is:

- Short-sleeved shirt
- Short pants
- Chemical resistant gloves of any waterproof material

Non-Agricultural Use Requirements

The requirements in this box apply to uses of this product that are NOT within the scope of the Worker Protection Standard for agricultural pesticides (40 CFR Part 170). The WPS applies when this product is used to produce agricultural plants on farms, forests, nurseries or greenhouses; do not allow people or pets to enter treatment area until the application is complete and the grass is dry.

INFORMATION

For use on newly seeded or over-seeded established plantings of the following grasses: Kentucky bluegrass, centipede grass, tall fescue, fine fescue (creeping red, chewing, hard), and perennial ryegrass.

Mesotrifluron may be applied to home lawns, recreational areas, golf courses and sod farms.

This product contains 0.15 pounds of mesotrifluron per acre per application. The maximum number of broadcast applications per site per year is three. Do not apply more than 0.45 pounds of mesotrifluron active ingredient per treatment site per year.

Apply on a calm day.

When making applications to newly seeded and over seeded established lawns and turf, apply mesotrifluron incorporated on fertilizer as the last step in the planting process. Best results will be obtained if 1/2 to 3/4 inch of water is provided by rainfall or irrigation within three days of treatment. Dry conditions following application may reduce herbicide activity.

Mesotrifluron may cause temporary whitening of turfgrass blades. In general, symptoms appear five to seven days after application and may last up to 14 days. Discoloration is most likely to occur on vigorous lawns and turf.

RESTRICTIONS

- Do not use this product on flowers, shrubs, trees, fruit or vegetable plants, Cans, dry fills and trees are especially susceptible to injury.
- Do not use on lawns that are being overseeded.
- Do not treat golf course greens with this product.
- Do not apply to Bentgrass, Bermudagrass (common), improved seeded Bermudagrass, Zoysiagrass, Deciduous, Kikuyagrass, Bahiagrass or Lippa lawns and turf.
- Do not apply an organophosphate or carbamate insecticide within seven days of an application of this product as turf injury may occur.

APPLICATION INSTRUCTIONS

Newly Seeded or Over Seeded Areas

Mesotrifluron applied as a starter fertilizer suppresses the emergence of crabgrass seedlings and controls broadleaf weeds without harming desirable turf grasses listed below. Turfgrass and lawns may be seeded and weeds controlled at the same time. Apply mesotrifluron incorporated on fertilizer as the last step in the planting process. **Spring Seeding:** For control of crabgrass, common chickweed, mouse ear chickweed, henbit, corn spurreck, hairy bittercress, leaf-alk, woodcock, other weed, volunteer ryegrass, sandolin, white clover, black medick, hop clover, timothy, ground ivy, broadleaf plantain, buckhorn plantain, Carolina planain, henbit, wildflower, wild radish, knotweed, and other broadleaf weeds, apply 187 pounds of fertilizer (0.15 pounds active ingredient per acre of mesotrifluron, pigweed, Kochia, wild radish, knotweed, and other broadleaf weeds, apply 187 pounds of fertilizer (0.15 pounds active ingredient per acre of mesotrifluron, pigweed, Kentucky bluegrass, centipede grass, tall fescue, creeping red, hard, and perennial ryegrass. Mesotrifluron may reduce the density of fine fescue seedlings. Mesotrifluron can be used on grass seed blends that contain less than 20% by weight of fine fescue (creeping red, chewing, hard).

For extended control of weeds listed above, a second application of 0.08% mesotrifluron on fertilizer can be applied six weeks after the first application. A second application will add in control of dandelion and perennials such as clover, ground ivy, and volunteer ryegrass. For extended season long control of crabgrass and foxtail, apply mesotrifluron in combination with a pre-emergence herbicide.

Fall Seeding: For control of crabgrass, common chickweed, mouse ear chickweed, henbit, corn spurreck, hairy bittercress, leaf-alk, knotweed, woodcock, dollar weed, volunteer ryegrass, sandolin, white clover, black medick, hop clover, timothy, ground ivy, broadleaf plantain, buckhorn plantain, Carolina planain, henbit, knotweed, wild radish, knotweed, and other broadleaf weeds, apply 187 pounds of fertilizer (0.15 pounds active ingredient per acre of mesotrifluron, pigweed, Kentucky bluegrass, centipede grass, tall fescue, creeping red, hard, and perennial ryegrass. Mesotrifluron may reduce the density of fine fescue seedlings. Mesotrifluron can be used on grass seed blends that contain less than 20% by weight of fine fescue (creeping red, chewing, hard).

WHAT TO EXPECT: Weed stems and leaves will begin to turn white within a few days of treatment. Weeds will turn brown and die. Two to three weeks may be required for complete control. Weeds not killed are often started and non-competitive in turf with flowering dandelion.

SUGGESTED SPREADER SETTINGS		4.3 #/1,000 sq. ft.	4.3 #/1,000 sq. ft.
Spreader		Leuco (patented)	1810
Lawnmower	4	ProGrass 30	1
Handheld	14	Syrter	4
Enduro (battery)	24	Syrter (pat model)	22
Leafy	40-1		

These settings were calibrated and field tested. However, age and condition of spreader, speed of operation and wetness of terrain may require slightly different settings for desired coverage. For best results, the product should be applied without slope or misfire. Determine the correct amount of product for the turf area to be treated. For best results apply half the rate in one direction and the remaining half rate at right angles to the first application. This procedure will minimize the chances for color streaking and ensure weed control.

STORAGE AND DISPOSAL

Do not contaminate water, food, or feed by storage or disposal. **PESTICIDE DISPOSAL:** Store in original package. Do not mix or add to other pesticides. Do not reuse or refill the container. Completely empty bag into application equipment. Then clean application equipment, including bag, in a sanitary, landfill or by incineration, or, by burning. Do not burn unless allowed by state or local authorities. If burned, stay out of smoke.

LIMITATION OF WARRANTY AND LIABILITY

Notice: Read This Limitation of Warranty and Liability Before Buying or Using This Product. If the Terms Are Not Acceptable, Return the Product at Once, Unopened, and the Purchase Price Will Be Refunded.

It is impossible to eliminate all risks associated with the use of this product. Such risks may arise from factors such as weather conditions, soil factors, off target movement, unconventional biotiny, presence of other materials, the manner of use or application, or other unknown factors, all of which are beyond the control of this company or Seller. These risks can cause: ineffectiveness of the product or other unknown factors; consequences. WHEN YOU BUY OR USE THIS PRODUCT, YOU AGREE TO ACCEPT THESE RISKS.

This company warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for the purpose stated in the Directions for Use, subject to inherent risks described above, when used in accordance with the Directions for Use under normal conditions. TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THIS COMPANY MAKES NO OTHER EXPRESS OR IMPLIED WARRANTY OF FITNESS OR MERCHANTABILITY OR ANY OTHER EXPRESS OR IMPLIED WARRANTY.

TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, NO EXCEPT SHALL THIS COMPANY OR SELLER BE LIABLE FOR ANY INCIDENTAL, CONSEQUENTIAL OR SPECIAL DAMAGES RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT TO THE EXTENT CONSISTENT WITH APPLICABLE LAW, THE EXCLUSIVE REMEDY OF THE USER OR BUYER AND THE EXCLUSIVE LIABILITY OF THIS COMPANY OR SELLER, FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY OR CONTRACT NEGLIGENCE, TORT, STRICT LIABILITY OR OTHER LEGAL THEORY) RESULTING FROM THE STORAGE, USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT, OR AT THE ELECTION OF THIS COMPANY OR SELLER, THE REPLACEMENT OF THE PRODUCT.

To the extent consistent with applicable law that allows such requirement, this company or its distributor must have prompt notice of any claim so that an immediate investigation of buyer's or user's claim can be made. Buyer and all users shall promptly notify this company or this company's distributor of any claims, whether based on contract, negligence, strict liability, other tort or otherwise or be barred from any remedy.


This Limitation of Warranty and Liability may not be amended by any oral or written agreement.

NOTICE: This product contains the secondary microconstituent Iron. Iron may stain concrete surfaces and should not be applied on dry or water dampened concrete and should be removed from these areas promptly by sweeping or blowing. Do not wash off with water.



Distributed by:
Lebanon Seaboard Corporation
 1600 E. Cumberland St. • Lebanon, PA 17042

Information regarding the contents and levels of metals in this product is available on the internet at:
<http://www.regulatory-info-lebesa.com>



ACCEPTED FOR REGISTRATION
 OCT 26 2017

New York State Department of Environmental Conservation
 Division of Materials Management
 Pesticide Product Registration
 23202-1 LB9611
 Doc ID: 554066

NET WEIGHT **40 LBS.** (18.14 KG)